



NAUSEA & VOMITING	
ADULT	PEDIATRIC
<b>BLS Procedures</b>	
Assess airway, suction airway as needed. Assess vitals. Obtain SpO2 Oxygen. Titrate to SpO2 to 94% or higher.	
<b>EMT Optional Scope</b>	
<b>ADMINISTRATION OF ONDANSETRON (ZOFRAN) BY EMT IS ONLY APPROVED FOR EMT'S THAT HAVE SUCCESSFULLY COMPLETED A STANISLAUS COUNTY EMS AGENCY APPROVED TRAINING.</b>	
<u><b>Adult Patients Only (18 Years and Up)</b></u>  <b>Ondansetron (Zofran) 4 mg PO</b> <ul style="list-style-type: none"> <li>May repeat once after 15 min</li> </ul>	<b>Pediatric Ondansetron (under 18 years old) not approved for EMT's</b>
<u><b>First Trimester Pregnancy (Less Than 12 Weeks)</b></u>  Withhold Ondansetron (Zofran) in First Trimester Pregnancy unless already using via prescription.	
<b>ALS Standing Orders</b>	
<b>IV access as needed.</b> <b>Obtain ECG.</b>	
<b>Ondansetron (Zofran) 4 mg PO/IM/IV/IO</b> <ul style="list-style-type: none"> <li>Slow IV push over 1 minute.</li> <li>Single max dose 4 mg.</li> <li>May repeat every 10 minutes.</li> <li>Total max dose 12 mg</li> </ul>	<u><b>2-3 Years Old</b></u>  <b>Ondansetron (Zofran) 2 mg IM/IV/IO</b> <ul style="list-style-type: none"> <li>Slow IV Push over 1 minute.</li> <li>No repeat</li> </ul>
<u><b>First Trimester Pregnancy (Less Than 12 Weeks)</b></u>  Withhold Ondansetron (Zofran) in First Trimester Pregnancy unless already using via prescription.	<u><b>4 Years and Older</b></u>  <b>Ondansetron (Zofran) 4 mg PO/IM/IV/IO</b> <ul style="list-style-type: none"> <li>Slow IV push over 1 minute.</li> <li>No repeat</li> </ul>
<b>Special Considerations</b>	
<ul style="list-style-type: none"> <li>Nausea and Vomiting are signs and symptoms of many other illnesses and or diseases.</li> <li>Have a high suspicion of cardiac related conditions or dysrhythmias.</li> </ul>	
<b>Base Hospital Orders Only</b>	
Contact Base Hospital Physician for Doses that exceed written protocol, or for patients under 2 years old.	